

## PAD VOLUME II

### References (First and Last Page)

1. Weitz JI, Byrne J, Clagett GP, et al. Diagnosis and treatment of chronic arterial insufficiency of the lower extremities: a critical review. *Circulation* 1996; 94:3026-49.
2. Ross R. Atherosclerosis-an inflammatory disease. *N Engl J Med* 1999; 340:115-26.
3. Vogt MT. Peripheral arterial disease: the epidemiology and burden of disease in older women. Continuing Medical Education (CME) Monograph on the Society for Vascular Medicine and Biology (SVMB) website. [Cited 2002]. No longer available.
4. McDermott MM, Fried L, Simonsick E, et al. Asymptomatic peripheral arterial disease is independently associated with impaired lower extremity functioning. *Circulation* 2000; 101: 1007-12.
5. McDermott MM, Greenland P, Liu K, et al. The ankle brachial index is associated with leg function and physical activity: the walking and leg circulation study. *Ann Intern Med* 2002; 136: 873-83.
6. Gerhard-Herman MD, Wittgen CM. Treatment and outcomes of peripheral atherosclerosis in women. CME Monograph on the SVMB website. [Cited 2002]. No longer available.
7. Ouriel K. Peripheral arterial disease. Seminar. *Lancet* 2001; 358: 1257-64.
8. Jaff MR. Severe peripheral arterial disease and critical limb ischemia: incidence, pathophysiology, presentation, methods of diagnosis. CME Monograph on the SVMB website. [Cited 2002]. No longer available.
9. Meijer WT, Hoes AW, Rutgers D, et al. Peripheral arterial disease in the elderly. The Rotterdam Study. *Arterioscler Thromb Vasc Biol* 1998; 18: 185-92.
10. Schainfeld RM. Management of peripheral arterial disease and intermittent claudication. *J Am Board Fam Pract* 2001; 14(6): 443-5.
11. TransAtlantic Inter-Society Consensus (TASC) Working Group. Epidemiology, natural history, risk factors. TASC document. *J Vasc Surg* 2000; 31(suppl): S23.
12. Sacco RL, Benjamin EJ, Broderick JP, et al. Public health burden of stroke. *Stroke* 1997; 28:1507-17.

280. Tsai AW, Folsom AR, Rosamond WD, Jones DW. Ankle-brachial index and 7-year ischemic stroke incidence. The ARIC study. *Stroke* 2001; 32: 1721-4.
281. Fowkes FG, Housley E, Cawood EH, et al. Edinburgh Artery Study: Prevalence of asymptomatic and symptomatic peripheral arterial disease in the general population. *Int J Epidemiol* 1991; 20: 384-92.
282. Leng GC, Fowkes FG, Lee AJ, et al. Use of ankle brachial pressure index to predict cardiovascular events and death: a cohort study. *BMJ* 1996; 313: 1440-3.
283. Stoffers HE, Rinkens PE, Kester AD, et al. The prevalence of asymptomatic and unrecognized peripheral arterial occlusive disease. *Int J Epidemiol* 1996; 25: 282-90.
284. Newman AB, Sutton-Tyrrell K, Kuller LH. Lower-extremity arterial disease in older hypertensive adults. *Arterioscler Thromb Vasc Biol* 1993; 13: 555-62.
285. Curb JD, Masaki K, Rodriguez B, et al. Peripheral artery disease and cardiovascular risk factors in the elderly. The Honolulu Heart Program. *Arterioscler Thromb Vasc Biol* 1996; 16: 1495-1500.
286. Bernstein EF, Dilley RF, Thomas WS, et al. Changing practice patterns in peripheral arterial disease. *Ann Vasc Surg* 1994; 8: 186-94.
287. De Vries SO, Donaldson MC, Hunink MGM. Contralateral symptoms after unilateral intervention for peripheral occlusive disease. *J Vasc Surg* 1998; 27: 414-21.
288. Tarry WC, Walsh DB, Birkmeyer NJO, et al. Fate of the contralateral limb after infrainguinal bypass. *J Vasc Surg* 1998; 27: 1039-48.
289. Collins TC, Petersen NJ, Suarez-Almazor M, Ashton CM. The prevalence of peripheral arterial disease in a racially diverse population. *Arch Intern Med* 2003; 163: 1469-1474.
290. Visser K, Idu MM, Buth J, et al. Duplex scan surveillance during the first year after infrainguinal autologous vein bypass grafting surgery: costs and clinical outcomes compared with other surveillance programs. *J Vasc Surg* 2001; 33: 123-30.
291. Wixon CL, Mills JL, Westerbrand A, et al. An economic appraisal of lower extremity bypass graft maintenance. *J Vasc Surg* 2000; 32: 1-12.