

## REFERENCES

(First and Last Pages)

1. Rowlands TE, Donnelly R. Medical therapy for intermittent claudication. *Eur J Vasc Endovasc Surg* 2007; 34: 314-21.
2. Hankey GJ, Norman PE, Eikebloom JW. Medical treatment of peripheral arterial disease. *J Am Med Assoc* 2006; 295: 547-53.
3. Balsano F, Coccheri S, Libretti A, et al. Ticlopidine in the treatment of intermittent claudication: a 21-month double blind trial. *J Lab Clin Med*. 1989; 114: 84-91.
4. Horsch S, Walther C. Ginkgo biloba special extract Egb 761 in the treatment of peripheral arterial occlusive disease (PAOD)—a review based on randomized, controlled studies. *Int J Clin Pharmacol Ther*. 2004; 42: 63-72.
5. Pittler MH, Ernst E. Ginkgo biloba extract for the treatment of intermittent claudication: a meta-analysis of randomized trials. *Am J Med*. 2000; 108:276-81.
6. Aronow WS, Nayak D, Woodworth S, Ahn C. Effect of simvastatin versus placebo on treadmill exercise time until the onset of intermittent claudication in older patients with peripheral arterial disease at six months and at one year after treatment. *Am J Cardiol*. 2003; 92: 711-712.
7. Mohler ER, Hiatt WR, Creager MA. Cholesterol reduction with atorvastatin improves walking distance in patients with peripheral arterial disease. *Circulation*. 2003; 108: 1481-86.
8. Pedersen TR, Kjekshus J, Pyörälä K, et al. Effect of simvastatin on ischemic signs and symptoms in the Scandinavian Simvastatin Survival Study (4S). *Am J Cardiol*. 1998; 81: 333-335.
9. Ahimastos AA, Lawler A, Reid CM, et al. Brief communication: ramipril markedly improves walking ability in patients with peripheral arterial disease. *Ann Intern med* 2006; 144: 660-4.
10. Thompson PD, Zimet R, Forbes WP, Zhang P. Meta-analysis of results from eight randomised, placebo-controlled trials on the effect of cilostazol on patients with intermittent claudication. *Am J Cardiol* 2002; 90: 1314-19.
11. IMS Healthcare. IMS MIDAS Database Search conducted 2007, November 28.
12. U.K. Medicines Information Pharmacists Group. [Internet.] Cilostazol. Treatment Alternatives. Available at <http://www.ukmi.nhs.uk/NewMaterial/html/docs/Cilostazol.pdf>.
13. Note for guidance on clinical investigation of medicinal products for the treatment of peripheral arterial occlusive disease. Committee for Proprietary Medical Products. The European Agency for the Evaluation of Medical Products. Available at <http://www.emea.europa.eu/pdfs/human/ewp/071498en.pdf>.

66. McDermott MM, Fried L, Simonsick E, et al. Asymptomatic peripheral arterial disease is independently associated with impaired lower extremity functioning. *Circulation* 2000; 101:1007-12.
67. McDermott MM, Greenland P, Liu K, et al. The ankle brachial index is associated with leg function and physical activity: the walking and leg circulation study. *Ann Intern Med* 2002; 136: 873-83.
68. Gardner AW, Montgomery PS. Impaired balance and higher prevalence of falls in subjects with intermittent claudication. *J Gerontol A Biol Sci Med Sci* 2001; 56(7): M454-8. Abstract.
69. Gardner AW, Montgomery PS. The relationship between history of falling and physical function in subjects with peripheral arterial disease. *Vasc Med* 2001; 6(4): 223-7. Abstract.
70. Guralnik JM, LaCroix AZ, Abbott RD, et al. Maintaining mobility in late life. I. Demographic characteristics and chronic conditions. *Am J Epidemiol* 1993; 137(8): 845-57.
71. Vogt MT, Cauley JA, Kuller LH, Nevitt MC. Functional status and mobility among elderly women with lower extremity arterial disease: the Study of Osteoporotic Fractures. *J Am Geriatr Soc* 1994; 42: 923-9. Abstract.
72. Criqui MH, Denenberg JO, Bird CE, et al. The correlation between symptoms and non-invasive test results in patients referred for peripheral arterial disease testing. *Vasc Med* 1996; 1:65-71. Abstract.
73. McDermott MM, Mehta S, Greenland P. Exertional leg symptoms other than intermittent claudication are common in peripheral arterial disease. *Arch Intern Med* 1999; 159(4): 387-92.

