

## **REFERENCES—First and Last Pages**

1. Prompers L, Huijberts M, Apelqvist J, et al. High prevalence of ischaemia, infection and serious comorbidity in patients with diabetic foot disease in Europe. Baseline results from the Eurodiale study. *Diabetologia* 2007; 50: 18-25.
2. Weitz JI, Byrne J, Clagett GP, et al. Diagnosis and treatment of chronic arterial insufficiency of the lower extremities: a critical review. *Circulation* 1996; 94: 3026-49.
3. Ross R. Atherosclerosis—an inflammatory disease. *N Engl J Med* 1999; 340: 115-26.
4. Criqui MH. Systematic atherosclerosis risk and the mandate for intervention in atherosclerotic peripheral arterial disease. *Am J Cardiol* 2001; 88(7B): 43J-7J.
5. Vogt MT. Peripheral arterial disease: the epidemiology and burden of disease in older women. Continuing Medical Education (CME) Monograph on the Society for Vascular Medicine and Biology (SVMB) website. [Cited 2002]. No longer available.
6. McDermott MM, Fried L, Simonsick E, et al. Asymptomatic peripheral arterial disease is independently associated with impaired lower extremity functioning. *Circulation* 2000; 101: 1007-12.
7. McDermott MM, Greenland P, Liu K, et al. The ankle brachial index is associated with leg function and physical activity: the walking and leg circulation study. *Ann Intern Med* 2002; 136: 873-83.
8. Nehler MR, Wolford H. Natural history and nonoperative treatment of chronic lower extremity ischemia. In: Rutherford, RB, editor. *Vascular surgery*. 6<sup>th</sup> ed. Philadelphia (PA): Elsevier Saunders; 2005. p.1083-94.
9. Biamino G, Scheinert D, Schmidt A, et al. Critical limb ischemia: new techniques for complex intervention. Malvern (PA): HMP Communications; 2004.
10. Garcia LA. Epidemiology and pathophysiology of lower extremity peripheral arterial disease. *J Endovasc Ther* 2006; 13(Suppl II): II-3-II-9.
11. Dormandy J, Heeck L, Vig S. Acute limb ischemia. *Seminars in Vascular Surgery* 1999; 12: 148-53.
12. Jaff MR, Biamino G. An overview of critical limb ischemia. *Endovascular Today* 2004; 3(2): 45-48.
13. Nanjundappa A, Laird JR. Critical limb ischemia: understanding the scope of the problem. *Endovascular Today* 2006; 5(7): 36-40.
14. Wolfe JHN, Wyatt MG. Critical and sub-critical ischaemia. *Eur J Vasc Endovasc Surg* 1997; 13: 578-82.
15. Holdsworth RJ, McCollum PT. Results and resource implications of treating end-stage limb ischaemia. *Eur J Vasc Endovasc Surg* 1997; 13: 164-73.

172. Williams DT, Harding KG, Price P. An evaluation of the efficacy of methods used in screening for lower-limb arterial disease in diabetes. *Diabetes Care* 2005; 28(9): 2206-10.
173. Goss DE, de Trafford J, Roberts VC, et al. Raised ankle/brachial pressure index in insulin-treated diabetic patients. *Diab Med* 1989; 6(7): 576-8. Abstract.
174. Silvestro A, Diehm N, Savolainen H, et al. Falsely high ankle-brachial index predicts major amputation in critical limb ischemia. *Vasc Med* 2006; 11(2): 69-74. Abstract.
175. Resnick HE, Lindsay RS, McDermott MM, et al. Relationship of high and low ankle brachial index to all-cause and cardiovascular disease mortality. The Strong Heart Study. *Circulation* 2004; 109: 733-9.
176. Wattanakit K, Folsom AR, Duprez DA, et al. Clinical significance of high ankle-brachial index: insights from the Atherosclerosis Risk in Communities (ARIC) Study. *Atherosclerosis* 2007; 190(2): 459-64.
177. Weatherly BD, Nelson JJ, Heiss G, et al. The association of the ankle-brachial index with incident coronary heart disease: the Atherosclerosis Risk in Communities (ARIC) study, 1987-2001. *BMC Cardiovasc Disord* 2007; 7: 3.
178. American Diabetes Association. Economic costs of diabetes in the U.S. in 2007. *Diabetes Care* 2008; 31: 596-615.
179. Bureau of Labor Statistics, Consumer Price Index, Archived Consumer Price Index Detailed Report Information, 2009 CPI Index Detailed Report Tables. Available at: [www.bls.gov/cpi/](http://www.bls.gov/cpi/). [Accessed 2010, September].