

REFERENCES (FIRST AND LAST PAGES)

1. Yost ML. Critical limb ischemia. United States epidemiology. Prevalence, market opportunities and analysis of the two most common comorbidities: diabetes and chronic kidney disease. Atlanta (GA): THE SAGE GROUP; 2010.
2. Wild S, Roglic G, Green A, et al. Global prevalence of diabetes. Estimates for the year 2000 and projections for 2030. *Diabetes Care* 2004; 27: 1047-53.
3. International Diabetes Federation. IDF Diabetes Atlas, 6th ed. Brussels, Belgium: International Diabetes Federation, 2013. <http://www.idf.org/diabetesatlas>.
4. The DECODE Study Group. Age- and sex-specific prevalence of diabetes and impaired glucose regulation in 13 European cohorts. *Diabetes Care* 2003; 26(1): 61-9.
5. Shelton N. Diabetes. In: Craig R, Mindell J, editors. Health Survey for England 2006. Volume I. Cardiovascular disease and risk factors in adults. United Kingdom: The Information Centre; 2008: p. 63-84. Available at: <http://www.ic.nhs.uk/pubs/hse06cvdandriskfactors>.
6. TransAtlantic Inter-Society Consensus (TASC) Working Group. Epidemiology, natural history, risk factors. *J Vasc Surg* 2000; 31(Suppl):S5-S34.
7. Yost ML. Critical limb ischemia Volume II. Western Europe epidemiology. Atlanta (GA): THE SAGE GROUP; 2010.
8. Yost ML. India peripheral arterial disease, thromboangiitis obliterans & critical limb ischemia epidemiology and markets. Atlanta (GA): THE SAGE GROUP; 2009.
9. Yost ML. South America: diabetes, peripheral artery disease and critical limb ischemia. Atlanta (GA): THE SAGE GROUP; 2013.
10. Weitz JI, Byrne J, Clagett GP, et al. Diagnosis and treatment of chronic arterial insufficiency of the lower extremities: a critical review. *Circulation* 1996; 94: 3026-49.
11. Ross R. Atherosclerosis-an inflammatory disease. *N Engl J Med* 1999; 340: 115-26.
12. Criqui MH. Systematic atherosclerosis risk and the mandate for intervention in atherosclerotic peripheral arterial disease. *Am J Cardiol* 2001; 88(7B): 43J-7J.
13. McDermott MM, Fried L, Simonsick E, et al. Asymptomatic peripheral arterial disease is independently associated with impaired lower extremity functioning. *Circulation* 2000; 101: 1007-12.
14. McDermott MM, Greenland P, Liu K, et al. The ankle brachial index is associated with leg function and physical activity: the walking and leg circulation study. *Ann Intern Med* 2002; 136: 873-83.

122. Hirsch AT, Criqui MH, Treat-Jacobsen D, et al. Peripheral arterial disease detection, awareness, and treatment in primary care. *JAMA* 2001; 286: 1317-24.
123. Zeymer U, Senges J, Parhofer KG, et al. Risk factors and event rates in patients with atherothrombotic disease in Germany: results of the German REACH registry. *Dtsch Arztebl Int* 2008; 105(45): 769-75.
124. Yost ML. Diabetic foot ulcers, peripheral artery disease and critical limb ischemia. Atlanta (GA): THE SAGE GROUP; 2010.
125. History of foot ulcer among persons with diabetes United States, 2000-2002. *Morb Mortal Wkly Rep* 2003 Nov 14: 52(45);1098-1102
126. Apelqvist J, Larsson J. What is the most effective way to reduce incidence of amputation in diabetic foot. *Diabetes Metab Res Rev* 2000; 16(Suppl 1): S75-S83.
127. Neil HA, Thompson AV, Thorogood M, et al. Diabetes in the elderly: the Oxford Community Diabetes Study. *Diabet Med* 1989; 6(7): 608-13. Abstract.
128. Walters DP. The distribution and severity of diabetic foot disease: a community study with comparison to a non-diabetic group. *Diabet Med* 1992; 9: 354-8.
129. Iversen MM, Midthjell K, Ostbye T, et al. History and factors associated with diabetic foot ulcers in Norway: the Nord-Trondelag Health Study. *Scand J Public Health* 2008; 36: 62-8.
130. Samann A, Tajiyeva N, Muller T, et al. Prevalence of diabetic foot syndrome at the primary care level in Germany: a cross-sectional study. *Diabet Med* 2008; 25: 557-63.
131. Behthel MA, Sloan FA, Belsky D, Feinglos MN. Longitudinal incidence and prevalence of adverse outcomes of diabetes mellitus in elderly patients. *Arch Intern Med* 2007; 167: 921-7.
132. Ogrin R, Miller J, Tennant J. Lessons to be learnt from data collection in a high risk foot clinic. *Primary Intervention* 2007; 15(3):96-014.
133. Palumbo, PJ, Melton LJ. Peripheral vascular disease and diabetes. In: *Diabetes in America*. p. 401-8.
134. Laclé A, Valero-Juan LF. Diabetes-related lower-extremity amputation incidence and risk factors: a prospective seven-year study in Costa Rica. *Rev Panam Salud Publica* 2012; 32(3): 192-8.
135. Yost ML. The cost of amputation in 2014. Lecture presented at the 4th annual Amputation Prevention Symposium. August 14-16, 2014 Palmer House Hilton, Chicago IL.