## THE COST OF CRITICAL LIMB ISCHEMIA

Mary L. Yost 404-520-6652 THE SAGE GROUP

## THE SAGE GROUP, LLC RESEARCH AND CONSULTING 23 Ridge Rd Beaufort SC 29907

Copyright Pending 2019

All rights reserved, including the right of reproduction in whole or in part in any form.

## **REFERENCES (FIRST AND LAST PAGES)**

- 1. Yost ML. The real cost of peripheral artery disease. Atlanta (GA): THE SAGE GROUP; 2011.
- 2. Mustapha JA, Katzen BT, Neville RF, et al. Determinants of long-term outcomes and costs in the management of critical limb ischemia: a population-based cohort study. J Am Heart Assoc. 2018 Aug 21;7(16):e009724.
- 3. Hartman M, Martin AB, Espinosa N, Catlin A, The National Health Expenditure Accounts Team. National health care spending In 2016: spending and enrollment growth slow after initial coverage expansions. Health Aff (Millwood). 2018 Jan;37(1):150-160.
- 4. Kolte D, Kennedy KF, Shishehbor MH, et al. Thirty-day readmissions after endovascular or surgical therapy for critical limb ischemia: analysis of the 2013 to 2014 Nationwide Readmissions Databases. Circulation. 2017 Jul 11;136(2):167-176.
- 5. Sambamoorthi U, Tan X, Deb A. Multiple chronic conditions and healthcare costs among adults. Expert Rev Pharmacoecon Outcomes Res. 2015;15(5):823-32.
- 6. The MetLife Mature Market Institute. The MetLife study of caregiving costs to working caregivers. 2011. [Internet. Accessed 2019, June.] Available at: https://www.caregiving.org/wp-content/uploads/2011/06/mmi-caregiving-costs-working-caregivers.pdf.
- 7. Dunbar SB, Khavjou OA, Bakas T, et al; American Heart Association. Projected costs of informal caregiving for cardiovascular disease: 2015 to 2035: A Policy Statement From the American Heart Association. Circulation. 2018 May 8;137(19):e558-e577.
- 8. Joo H, Zhang P, Wang G. Cost of informal care for patients with cardiovascular disease or diabetes: current evidence and research challenges. Qual Life Res. 2017 Jun;26(6):1379-1386.
- 9. Dua A, Desai SS, Patel B, et al. Preventable complications driving rising costs in management of patients with critical limb ischemia. Ann Vasc Surg. 2016 May;33:144-8.
- 10. Agarwal S, Sud K, Shishehbor MH. Nationwide trends of hospital admission and outcomes among critical limb ischemia patients: from 2003-2011. J Am Coll Cardiol. 2016 Apr 26; 67(16):1901-13.
- 11. Martinez RA, Shnayder M, Parreco J, et al. Nationally representative readmission factors in patients with claudication and critical limb ischemia. Ann Vasc Surg. 2018 Oct;52:96-107.
- 12. Yost ML. Critical limb ischemia by Rutherford Category: prevalence and markets in patients and limbs. Beaufort (SC); THE SAGE GROUP; 2017.

- 159. Fingar K. (Truven Health Analytics), Washington R (AHRQ). Trends in Hospital Readmissions for Four High-Volume Conditions, 2009–2013. HCUP Statistical Brief #196. November 2015. Agency for Healthcare Research and Quality, Rockville, MD. [Internet. Accessed 2019, May.] Available at: <a href="http://www.hcup-us.ahrq.gov/reports/statbriefs/sb196-Readmissions-Trends-High-Volume-Conditions.pdf">http://www.hcup-us.ahrq.gov/reports/statbriefs/sb196-Readmissions-Trends-High-Volume-Conditions.pdf</a>.
- 160. Vogel TR, Kruse RL. Risk factors for readmission after lower extremity procedures for peripheral artery disease. J Vasc Surg. 2013 Jul;58(1):90-7.e1-4.
- 161. Jones CE, Richman JS, Chu DI, et al. Readmission rates after lower extremity bypass vary significantly by surgical indication. J Vasc Surg. 2016 Aug;64(2):458-464.
- 162. Bodewes TC, Soden PA, Ultee KH, et al. Risk factors for 30-day unplanned readmission following infrainguinal endovascular interventions. J Vasc Surg. 2017;65:484–494.e3.
- 163. Reed GW, Raeisi-Giglou P, Kafa R, et al. Hospital readmissions following endovascular therapy for critical limb ischemia: associations with wound healing, major adverse limb events, and mortality. J Am Heart Assoc. 2016;5:e003168.
- 164. Orr NT, El-Maraghi S, Korosec RL, et al. Cost analysis of vascular readmissions after common vascular procedures. J Vasc Surg. 2015 Nov;62(5):1281-7.e1.
- 165. Prompers L, Schaper N, Apelqvist J, et al. Prediction of outcome in individuals with diabetic foot ulcers: focus on the differences between individuals with and without peripheral arterial disease: the EURODIALE Study. Diabetologia. 2008;51:747–755.
- 166. Lavery LA, Armstrong DG, Murdoch DP, et al. Validation of the Infectious Diseases Society of America's diabetic foot infection classification system. Clin Infect Dis. 2007;44:562–565.
- 167. Melvin JC, Smith JB, Kruse RL, Vogel TR. Risk factors for 30-day hospital re-admission with an infectious complication after lower-extremity vascular procedures. Surg Infect (Larchmt). 2017 Apr;18(3):319-326.
- 168. Looser PM, Feldman DN. Thirty-day readmissions for critical limb ischemia: ready for a new quality metric? Circulation. 2017;136:177–179.
- 169. Brooke BS, Kraiss LW, Stone DH, et al. Improving outcomes for diabetic patients undergoing revascularization for critical limb ischemia: does the quality of outpatient diabetic care matter? Ann Vasc Surg. 2014 Oct;28(7):1719-28.

## **CONTACT INFORMATION**

Mary L. Yost President Telephone (404) 520-6652 yost@thesagegroup.us